## **EUROPEAN PATENT OFFICE**



## Patent Abstracts of Japan

PUBLICATION NUMBER

: 09009903

PUBLICATION DATE

: 14-01-97

APPLICATION DATE

: 30-06-95

APPLICATION NUMBER

: 07165461

APPLICANT: FUJITSUKO KK;

INVENTOR: OKUDAIRA TAKENORI;

INT.CL.

: A23L 1/20 C12N 1/20 //(C12N 1/20 , C12R 1:125 )

TITLE

FERMENTED SOYBEAN FUNGUS AND PRODUCTION OF FERMENTED SOYBEAN

WITH THE SAME AND FERMENTED SOYBEAN

ABSTRACT :

PURPOSE: To obtain a fermented soybean fungus belonging to Bacillus subtilis, having a hemicellulose-decomposing ability, and capable of producing fermented soybeans in which the seed coats of the soybeans are partially decomposed to improve the digestive

absorbability.

CONSTITUTION: A fermented soybean fungus comprises Bacillus subtilis neomild 710 (FERM P-14843) belonging to the genus Bacillus subtilis, having a hemicellulose-decomposing ability or an ability to decompose the hemicellulose of soybean seed coats with xylan, a xylanase activity of ≥10 units and a threading ability of ≥50cm on a GSP medium. Fermented soybeans are preferably produced with the fermented soybean fungus.

COPYRIGHT: (C)1997,JPO

	•
	•

Publication number:

**0 287 699** 

(2

### **EUROPEAN PATENT APPLICATION**

Application number: 87106255.0

<sup>(22)</sup> Date of filing: 29.04.87

⑤ Int. Cl.<sup>4</sup> C12N 1/20 , A23K 1/00 , A23K 1/18 , //(C12N1/20, C12R1:00)

The title of the invention has been amended (Guidelines for Examination in the EPO, A-III, 7.3).

The microorganism(s) has (have) been deposited with Fermentation Research Institute under number(s) FERM BP-1096.

- Priority: 25.02.87 JP 40097/87
- 43 Date of publication of application: 26.10.88 Bulletin 88/43
- Designated Contracting States:
  DE FR GB NL

- Applicant: The Calpis Food Industry Co., Ltd. 20-3, 2-chome Ebisu-Nishi Shibuya-ku Tokyo(JP)
- inventor: Iwanami, Takefumi
  5-4-2, Kurikidai Asau-ku
  Kawasaki-shi Kanagawa-ken(JP)
  Inventor: Maruta, Kiyoshi
  Senchurihaitsu-Machida 6-105 1-4-8,
  Minami-Naruse
  Machida-shi Tokyo(JP)
  Inventor: Murota, Ichiya
  1-3-16, Sagamihara Sagamihara-shi
  Kanagawa-ken(JP)
  Inventor: Miyazaki, Hiroshi
  2-22-11, Kamishakujii
  Nerima-ku Tokyo(JP)
- Representative: Thomsen, Dieter, Dr.rer.nat.
  Kaiser-Ludwig-Platz 6
  D-8000 München 2(DE)
- Feeds containing bacillus subtilis C-3102.

By giving mammals, fowls, fish, etc. feeds containing <u>Bacillus</u> <u>subtilis</u> C-3102 (FERM BP-1096), an excellent body weight gain and feed efficiency can be obtained.

EP 0 287 699 A2

Xerox Copy Centre

## FEEDS

### Field of Industrial Application

The present invetion relates to feeds containing Bacillus subtilis C-3102.

Bacillus subtilis C-3102 in accordance with the present invention exhibits remarkable effects of increasing weight and improving feed efficiency, when it is given to animals, such as mammals, fowls, and fishes as a feedstuff.

Accordingly, the present invention plays an important role not only in the art of microorganisms but also in the arts of livestock indusry and fisheries.

### Prior Art

15

Animal meat is very important as a foodstuff or a feedstuff. Accordingly, it has been strongly desired in the industrial world to efficiently raise economic animals.

For this reason, extensive investigations have been made from old on a formulation ratio of main ingredients which are inexpensive and have good nutrition balance, vitamins and minerals as trace components, and the like. Further a variety of probiotics (live bacterial agents) have been examined and the effects thereof have been noted to a certain extent but are yet unsatisfactory.

The present invention relates to feeds containing <u>Bacillus</u> <u>subtilis</u> C-3102; such bacteria have been neither found heretofore nor has it been known that this new strain can show an excellent effect as a probiotic.

25

30

### Problem to be solved by the Invention

As described above, it is the actual situation that any satisfactory probiotic having excellent safety and having a weight-increasing effect is unknown in the current feed industry and microorganism industry.

### Means for solving the Problem

35

In view of the circumstances described above, the present invention has been made and from the viewpoint that there is no alternative other than finding hitherto unknown, new bacteria in order to solve the problem described above, various screening tests have been repeated. As a result, a novel strain belonging to the genus <u>Bacillus</u> has been discovered and has been named <u>Bacillus subtilis</u> C-3102.

A useful and novel finding has been noted that when this <u>Bacillus subtilis</u> C-3102 is incorporated and contained in animal feeds, it can accelerate the gain of weight in animals and at the same time, can improve feed efficiency; and as a result of further investigations, the present invention has been accomplished.

Namely, the present invention relates to probiotics comprising <u>Bacillus</u> <u>subtilis</u> C-3102 and relates to feeds containing Bacillus subtilis C-3102.

The strain used in the present invention is <u>Bacillus subtilis</u> C-3102 newly isolated from the natural world by the present inventors.

1

Bacillus subtilis C-3102 has been deposited in Fermentation Research Institute, Agency of Industrial Science and Technology, Japan, under FERM BP-1096. Bacteriological properties of this strain are as follows.

### A. Morphological properties

- (1) Cells: rods having a width of 0.6 to 1.0  $\mu m$  and a length of 1.5 to 2.0  $\mu m$ .
- (2) Gram-positive
- (3) Spores: oval
- (4) Mobility: positive
- (5) Aerobic

10

5

- B. Growth Conditions in Various Media
  - (1) Nutrient agar: excellent
  - (2) Anaerobic glucose bouillon: -
  - (3) 7 % NaCl: +
- C. Physiological Properties

20

25

30

35

45

50

- (1) Growth at 50 °C:+
- (2) Growth at pH 5.7:+
- (3) Utilization of citrate: +
- (4) Acid production from carbohydrates arabinose, glucose, xylose, mannitol: +
  - (5) VP reaction: +
  - (6) Hydrolysis of starch: +
  - (7) Reduction of nitrate: +
  - (8) Production of indole: -
  - (9) Liquefaction of gelatin: +
  - (10) Decomposition of casein: +
  - (11) Film formation on liquid medium: + \*
  - (12) Coagulation of milk: -
  - (13) Peptonization of milk: +
  - (14) Catalase: +

Next, a method for culturing Bacillus subtilis C-3102 of the present invention will be described.

As media, there can be widely used media ordinarily used for culturing microorganisms containing carbon sources, nitrogen sources, inorganic matters, vitamins, amino acids, etc. The carbon source can be any carbon compounds which are assimilable and, for example, glucose, sucrose, starch, molasses, etc. 40 can be employed. The nitrogen source can be any nitrogen compounds which are utilizable and, for example, peptone, meat extract, casein acid hydrolysate, ammonium sulfate, etc. can be employed. In addition, phosphates, salts of metals such as magnesium, sodium, potassium, calcium, iron, manganese, etc.; vitamins, amino acids, defoaming agents, surfactants, etc. can also be used, depending upon necessity.

As the media, both liquid media and solid media can be used and incubation under aerobic conditions is appropriate.

Incubation can be suitably effected at an initial pH of the medium is pH of 5 to 9, preferably 6 to 8, at culturing temperatures at 20 to 50 °C, preferably 35 to 40°C and for a culturing time period of 12 hours to 7

The thus obtained culture can be used as it is or as washed bacteria. Further the culture or bacteria can also be used as it is or after additives (carries innoxious to animals) are added thereto followed by drying or preparations thereof are produced. The probiotics of the present invention include these forms for use, which comprises Bacillus subtilis C-3102. It is sufficient that Bacillus subtilis C-3102 be incorporated in an amount of 10 to 1012 cells, preferably 104 to 108 cells, per 1 g of a feed but the amount is not limited only 55 to such a range.

When the feed containing this bacteria is given to animals, a great weight-increasing effect can be obtained and feed conversion can also be greatly improved.

Moreover, the feed containing the bacteria is extremely effective for a variety of animals, in addition to

mammals such as cows, pigs, horses, sheep, etc., various fowls such as chicken, quails, etc., and fishes. Furthemore, the feed is also useful for various pet animals such as dogs, cats, goldfishes, canaries, etc.

Hereafter preparation examples of the bacteria and reference examples showing fatting tests using the obtained probiotic will be described in detail.

Preparation Example 1

5

Medium obtained by dissolving 400 g of soybean peptone, 10 g of dibasic potassium phosphate and 200 g of molasses in 60 liters of tap water and adjusting pH to 7.5 with 1 N sodium hydroxide solution was charged in a jar fermenter and sterilized at 121°C for 15 minutes. A culture solution of Bacillus subtilis C-3102, FERM BP-1096, which had been previously subjected to preliminary incubation, was inoculated thereinto followed by spinner culture under aeration at 37°C for 40 hours.

The thus obtained culture solution was centrifuged to collect the cells. After drying, the cells was incorporated in skim milk powder to give 10 kg of a probiotic of <u>Bacillus subtilis</u> C-3102. The viable count contained in this probiotic was 1 x 10<sup>s</sup> cells/g.

### 20 Preparation Example 2

To 5 kg of soybean oil meal granules (Fujinic Ace 500, manufactured by Fuji Purina Protein Co., Ltd.) was added 5 kg of tap water followed by sterilization at 121°C for 120 minutes. A culture solution of Bacillus subtilis C-3102, FERM BP-1096, which had been previously subjected to preliminary incubation, was inoculated thereinto followed by culture at 37°C for 40 hours.

The thus obtained culture solution was dried and reduced into powder. The powder was incorporated in calcium carbonate to give 4 kg of a probiotic of Bacillus subtilis C-3102. The viable count contained in this probiotic was  $1 \times 10^{10}$  cells/g.

#### Example 1

30

40

45

50

Male ICR mice of 6 age in weeks were divided into 2 groups (test gr. and control gr.) of 10 each and the feeds having the compositions as shown in Table 1 were given to the respective groups to conduct growth test. The probiotic obtained in Preparation Example 1 by incorporating into skim milk powder was used. The viable count in the feed in which the probiotic was added was 1 x 10<sup>6</sup> cells/g. The results are as shown in Table 2.

(unit: wt%)

29.0

33.6

4.6

[100]

103.0

22.4

Table 1

5		Invention	Control
10	CE-2(feed for mouse, manufactured by CLEA JAPAN, INC.	99.9	99.9
	Skim milk powder	-	0.1
15	Probiotic of the present invention	0.1	-
20	Table 2	<b>.</b>	
		Invention	Control
25	Feeding period (days)	19	19

Mean body weight at start(g)

Mean body weight at end(g)

Mean body weight gain(g)

[Body weight gain index]

Mean feed intake(g)

Feed conversion ratio

As shown in Table 2, it is understood that by feeding the formula feed added with the probiotic of the present invention, an outstanding body weight-increasing effect was obtained and at the same time, the feed conversion ratio was also improved.

29.0

35.2

6.2

[135]

109.2

17.6

### 45 Example 2

Piglings of 3 to 4 age in weeks were divided into 2 groups (test gr. and control gr.) of 6 each and the feeds having the compositions as shown in Table 3 were given to the respective groups to conduct growth test. The probiotic obtained in Preparation Example 2 by incorporating into calcium carbonate was used. The viable count in the feed in which the probiotic was added was 1 x 10<sup>6</sup> cells/g. The resuls are as shown in Table 4.

55

30

Table 3		(unit: wt%)	
		Invention	Control
5	Bakery waste (PANBIS, manufactured by Koike-shoten Co.)	37	37
10	Skim milk powder	35.99	35.99
.0	Feed which has adsorbed soybean extract (BITAZE 8, manufactured by Enaze Sangyo Co., Ltd.)	11	11
15	G]ucose	6	6
20	Dehulled soybean powder (PAFESU 109, manufactured by Kikkoman Co., Ltd.)	3	3
0.5	Vitamins, minerals and other additives	7	7
25	Calcium carbonate	-	0.01
30	Probiotic of the present invention	0.01	-

Table 4

5		Invention	Control
	Feeding period (days)	21	21
	Mean body weight at start(kg)	6.1	6.1
10	Mean body weight at end (kg)	13.9	13.2
	Mean body weight gain(kg)	7.8	7.1
15	[Body weight gain index]	[110]	[100]
	Mean feed intake (kg)	67.9	67.3
20	Feed conversion ratio	1.45	1.58
	Observation of feces (points: cf. Note)	19	54

Note: Total points of each group when normal feces, soft feces, diarrhea and watery feces were designated 0, 1, 2 and 3 points, respectively.

As shown in Table 4, it is understood that by feeding the formula feed added with the probiotic of the present invention, an outstanding body weight-increasing effect was obtained and at the same time, the feed conversion ratio was also improved, and further that the occurrence of diarrhea and soft feces was greatly reduced.

### Example 3

40

25

30

Broilers of 2 to 3 age in days were divided into 2 groups (test gr. and control gr.) of 30 each (15 females and 15 males) and the feeds having the compositions as shown in Table 5 were given to the respective groups to conduct growth test.

The probiotic obtained in Preparation Example 1 by incorporating into skim milk powder was used. The viable count in the feed in which the probiotic was added was  $1 \times 10^6$  cells.g. The results are as shown in Table 6.

50

Table 5 (unit: wt%)

5		Invention	Control
	Commercial formula feeds for broilers for use in a former feeding period (20 days) and for use in a latter feeding period (36 days) (manufactured by Kumiai Feed Co.)	99.9	99.9
15	Skim milk powder	-	0.1
	Probiotic of the present invention	0.1	-

Table 6

25		Invention	Control
	Feeding period (days) Mean body weight at start(kg)	56 40	56 40
30	Mean body weight at end(kg)	2150	2000
	Mean body weight gain(g)	2110	1960
35	[Body weight gain index]	[108]	[100]
	Mean feed intake (kg)	4642	4528
	Feed conversion ratio	.2.20	2.31

As shown in Table 6, it is understood that by feeding the formula feed added with the probiotic of the present invention, an outstanding body weight-increasing effect was obtained and at the same time, the feed conversion ratio was also improved.

Example 4

Chicken for harvesting eggs were divided into 2 groups (test gr. and control gr.) of 30 each and the feeds having the compositions shown in Table 7 were given to the respective groups to conduct a test. The probiotic obtained in Preparation Example 1 by incorporating into skim milk powder was used. The viable count in the feed in which the probiotic was added was 1 x 10<sup>6</sup> cells/g. The results are as shown in Table 8.

55

40

45

(unit: wt%)

0.1

Commercial formula 99.9 99.9 feed (Kumiai Feed Co.)

Probiotic of the present invention 0.1

Table 8

Skim milk powder

Table 7

.²C		Invention	Control
	Feeding period (days)	28	28
	Total number of laid eggs	712	672
25	[Laid-egg number index]	[106]	[100]
	Mean egg weight (g)	60.2	59.2
30	[Egg weight index]	[102]	[100]

As shown in Table 8, it is understood that by feeding the formura feed added with the probiotic of the present invention, the number of laid eggs increased and at the same time, the mean egg weight also increased.

### Example 5

Rainbow trout fry of approximately 14 g were divided into 2 groups (test gr. and control gr.) of 30 each and the feeds having the compositions as shown in Table 9 were given to the respective groups to conduct growth test. The probiotic obtained in Preparation Example 1 by incorporating into skim milk powder was used. The viable count in the feed in which the probiotic was added was 1 x 10<sup>6</sup> cells/g. The results are as shown in Table 10.

50

10

15

	Table 9	(unit: wt%)	
		Invention	Control
5	Northern-sea fish meal	55	55
	α-Starch	20	20
10	Dextrin	5	5
	Oil mixture	10	10
15	Cellulose	3	3
	Vitamins, minerals and other additives	6.9	6.9
20	Skim milk powder	-	0.1
25	Probiotic of the present invention	0.1	-
25	Table 10		
		Invention	Control
30	Feeding period (days)	45	45
	Mean body weight at start(g)	14.3	14.0
35	Mean body weight at end (g)	39.6	37.6
	Mean body weight gain(g)	25.3	23.6
40	[Body weight gain index]	[107]	. [100]

As shown in Table 10, by feeding the formula feed added with the probiotic of the present invention, an outstanding body weight-increasing effect can be noted.

### Example 6

Calves of 1 week age after birth were divided into 2 groups (test gr. and control gr.) of 4 each and the feeds having the compositions as shown in Table 11 were given to the respective groups to conduct growth test. The probiotic obtained in Preparation Example 2 by incorporating into calcium carbonate was used. The viable count in the feed in which the probiotic was added was 1  $\times$  10<sup>6</sup> cells/g. Upon peforming the test, 1 part by weight of the feed was dissolved and dispersed in 7 parts by weight of hot water, which was then fed to the calves. The results are as shown in Table 12.

. .

	Table 11	(unit	: wt%)
		Invention	Control
5	Skim milk powder	70	70
	Whey	10	10
10	Beef tallow	12.49	12.49
	Lecithin	0.5	0.5
15	Vitamins, minerals and other additives	7	7
	Calcium carbonate	-	0.01
20	Probiotic of the present invention	0.01	-
25	Table 12		
20		Invention	Control
30	Feeding period (days)	28	28
	Mean body weight at start(kg)	43.7	43.9
35	Mean body weight at end(kg)	75.3	73.9
	Mean body weigt gain (kg)	31.6	30
40	[Body weight gain index]	[105]	[100]
	Mean feed intake (kg)	44.9	44.1
<b>4</b> 5	Feed conversion ratio	1.42	1.47
	Observation of feces (points: cf. Note)	29	45
50	Note: Total points of each group	when normal feces	, soft feces,
	diarrhea and watery feces were d	esignated 0, 1, 2	and 3 points,
55	respectively.		

# Claims 1. A probiotic for animals which comprises Bacillus subtilis C-3102 (FERM BP-1096) having bacteriological properties described below: (1) Gram-positive, (2) Cells: rods, (3) Spores: oval, 10 (4) Mobility: positive, (5) Aerobic, (6) Catalase: positive, (7) Growth at 50°C: +, (8) Growth at pH 5.7: +, 15 (9) Utilization of citrate: +, (10) Acid production from saccharides: arabinose, glucose, xylose, mannitol: +. (11) VP reaction: +, (12) Hydrolysis of starch: +, 20 (13) Reduction of nitrate: +, (14) Production of indole: -, (15) Liquefaction of gelatin: +, (16) Decomposition of casein: +. (17) Film formation on liquid medium: +, 25 (18) Coagulation of milk: -, and (19) Peptonization of milk: +, 2. A feed containing Bacillus subtilis C-3102 (FERM BP-1096) as defined in claim 1. 30 35 40 **4**5

13

50

		•

1 Publication number:

**0 287 699** A3

(12)

## **EUROPEAN PATENT APPLICATION**

- 2) Application number: 87106255.0
- 22 Date of filing: 29.04.87

(5) Int. Cl.4: C12N 1/20 , A23K 1/00 , A23K 1/18 , //(C12N1/20, C12R1:00)

- Priority: 25.02.87 JP 40097/87
- Date of publication of application: 26.10.88 Bulletin 88/43
- Designated Contracting States:
  DE FR GB NL
- Date of deferred publication of the search report: 26.07.89 Bulletin 89/30
- Applicant: The Calpis Food Industry Co., Ltd. 20-3, 2-chome Ebisu-Nishi Shibuya-ku Tokyo(JP)
- 2 Inventor: Iwanami, Takefumi
  5-4-2, Kurikidai Asau-ku
  Kawasaki-shi Kanagawa-ken(JP)
  Inventor: Maruta, Kiyoshi
  Senchurihaitsu-Machida 6-105 1-4-8,
  Minami-Naruse
  Machida-shi Tokyo(JP)
  Inventor: Murota, Ichiya
  1-3-16, Sagamihara Sagamihara-shi
  Kanagawa-ken(JP)
  Inventor: Miyazaki, Hiroshi
  2-22-11, Kamishakujii
  Nerima-ku Tokyo(JP)
- Representative: Thomsen, Dieter, Dr.rer.nat. Kaiser-Ludwig-Platz 6 D-8000 München 2(DE)
- Feeds containing bacillus subtilis C-3102.
- ⑤ By giving mammals, fowls, fish, etc. feeds containing Bacillus subtilis C-3102 (FERM BP-1096), an excellent body weight gain and feed efficiency can be obtained.

P 0 287 699 A

## **EUROPEAN SEARCH REPORT**

Application Number

EP 87 10 6255

]	DOCUMENTS CONSIL	DERED TO BE RELEVA	ANT	
ategory	Citation of document with inc of relevant pass		Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Ct. 4)
A	GB-A-1 069 400 (NAG * Claims 1-8 *	ASE & CO., LTD)	1,2	C 12 N 1/20 A 23 K 1/00
A	CHEMICAL ABSTRACTS, March 1987, page 493 65874q, Columbus, Oh 905 (V. BASTL et al. * Abstract *	, ab2tract no. io, US; & CS-A-232	1,2	A 23 K 1/18 // (C 12 N 1/20 C 12 R 1:00)
A	US-A-2 132 621 (A.A * Claim 1 *	. HORVATH)	1,2	
A	PATENT ABSTRACTS OF 251 (C-252)[1688], 1 & JP-A-59 128 334 (S KENKYUSHO K.K.) 24-0	.6th November 1984; SANWA KAGAKU	1,2	
				TECHNICAL FIELDS SEARCHED (Int. Cl.4)
				A 23 K
				C 12 N C 12 R
	The present search report has b	een drawn up for all claims		
711	Place of search	Date of completion of the scar	1	Examiner
. H	E HAGUE	26-04-1989	DEK	EIREL M.J.
Y:pa du	CATEGORY OF CITED DOCUMES  inticularly relevant if taken alone articularly relevant if combined with and acument of the same category chnological background	E: earlier pat after the f other D: document L: document	cited in the application	blished on, or on
O: 0	on-written disclosure termediate document		f the same patent fam	